

European Evidence-based (S3) Guideline for the Treatment of Acne - Update 2016

Evaluated studies:

- [1] Adhikary J, Satyanarayana V, Bhandare B, Vivekananda. A comparative study to evaluate the efficacy and safety of topical isotretinoin and clindamycin versus adapalene and clindamycin in the treatment of grade I - II acne vulgaris of face. International Journal of Pharmaceutical Sciences Review and Research. 2014;29: 175-179.
- [2] Al-Mishari MA. Clinical and bacteriological evaluation of tetracycline and erythromycin in acne vulgaris. Clin Ther. 1987;9; 273-280.
- [3] Alirezai M, Gerlach B, Horvath A, Forsea D, Briantais P, Guyomar M. Results of a randomised, multicentre study comparing a new water-based gel of clindamycin 1% versus clindamycin 1% topical solution in the treatment of acne vulgaris. Eur J Dermatol. 2005;15; 274-278.
- [4] Babaeenejad SH, Fouladi RF. The efficacy, safety, and tolerability of adapalene versus benzoyl peroxide in the treatment of mild acne vulgaris: a randomized trial. J Drugs Dermatol. 2013;12; 790-794.
- [5] Becker LE, Bergstresser PR, Whiting DA, Clendenning WE, Dobson RL, Jordan WP, et al. Topical clindamycin therapy for acne vulgaris. A cooperative clinical study. Arch Dermatol. 1981;117; 482-485.
- [6] Berger R, Barba A, Fleischer A, Leyden JJ, Lucky A, Pariser D, et al. A double-blinded, randomized, vehicle-controlled, multicenter, parallel-group study to assess the safety and efficacy of tretinoin gel microsphere 0.04% in the treatment of acne vulgaris in adults. Cutis. 2007;80; 152-157.
- [7] Berger R, Rizer R, Barba A, Wilson D, Stewart D, Grossman R, et al. Tretinoin gel microspheres 0.04% versus 0.1% in adolescents and adults with mild to moderate acne vulgaris: a 12-week, multicenter, randomized, double-blind, parallel-group, phase IV trial. Clin Ther. 2007;29; 1086-1097.
- [8] Bhavsar B, Choksi B, Sanmukhani J, Dogra A, Haq R, Mehta S, et al. Clindamycin 1% Nano-emulsion Gel Formulation for the Treatment of Acne Vulgaris: Results of a Randomized, Active Controlled, Multicentre, Phase IV Clinical Trial. J Clin Diagn Res. 2014;8; YC05-09.
- [9] Bleeker J. Tolerance and efficacy of erythromycin stearate tablets versus enteric-coated erythromycin base capsules in the treatment of patients with acne vulgaris. J Int Med Res. 1983;11; 38-41.
- [10] Bleeker J, Hellgren L, Vincent J. Effect of systemic erythromycin stearate on the inflammatory lesions and skin surface fatty acids in acne vulgaris. Dermatologica. 1981;162; 342-349.
- [11] Bojar RA, Eady EA, Jones CE, Cunliffe WJ, Holland KT. Inhibition of erythromycin-resistant propionibacteria on the skin of acne patients by topical erythromycin with and without zinc. Br J Dermatol. 1994;130; 329-336.
- [12] Borglund E, Hagermark O, Nord CE. Impact of topical clindamycin and systemic tetracycline on the skin and colon microflora in patients with acne vulgaris. Scand J Infect Dis Suppl. 1984;43; 76-81.
- [13] Bossuyt L, Bosschaert J, Richert B, Cromphaut P, Mitchell T, Al Abadie M, et al. Lymecycline in the treatment of acne: an efficacious, safe and cost-effective alternative to minocycline. Eur J Dermatol. 2003;13; 130-135.
- [14] Bowman S, Gold M, Nasir A, Vamvakias G. Comparison of clindamycin/benzoyl peroxide, tretinoin plus clindamycin, and the combination of clindamycin/benzoyl peroxide and tretinoin plus clindamycin in the treatment of acne vulgaris: a randomized, blinded study. J Drugs Dermatol. 2005;4; 611-618.
- [15] Braathen LR. Topical clindamycin versus oral tetracycline and placebo in acne vulgaris. Scand J Infect Dis Suppl. 1984;43; 71-75.
- [16] Bucknall JH, Murdoch PN. Comparison of tretinoin solution and benzoyl peroxide lotion in the treatment of acne vulgaris. Curr Med Res Opin. 1977;5; 266-268.

- [17] Callender VD, Young CM, Kindred C, Taylor SC. Efficacy and Safety of Clindamycin Phosphate 1.2% and Tretinoin 0.025% Gel for the Treatment of Acne and Acne-induced Post-inflammatory Hyperpigmentation in Patients with Skin of Color. *J Clin Aesthet Dermatol.* 2012;5; 25-32.
- [18] Cambazard F. Clinical efficacy of Velac, a new tretinoin and clindamycin phosphate gel in acne vulgaris. *J Eur Acad Dermatol Venereol.* 1998;11 Suppl 1; S20-27; discussion S28-29.
- [19] Cavicchini S, Caputo R. Long-term treatment of acne with 20% azelaic acid cream. *Acta Derm Venereol Suppl (Stockh).* 1989;143; 40-44.
- [20] Chalker DK, Lesher JL, Jr., Smith JG, Jr., Klauda HC, Pochi PE, Jacoby WS, et al. Efficacy of topical isotretinoin 0.05% gel in acne vulgaris: results of a multicenter, double-blind investigation. *J Am Acad Dermatol.* 1987;17; 251-254.
- [21] Chandrashekha BS, Anitha M, Ruparelia M, Vaidya P, Aamir R, Shah S, et al. Tretinoin nanogel 0.025% versus conventional gel 0.025% in patients with acne vulgaris: A randomized, active controlled, multicentre, parallel group, phase iv clinical trial. *Journal of Clinical and Diagnostic Research.* 2015;9; WC04-WC09.
- [22] Choudhury S, Chatterjee S, Sarkar DK, Dutta RN. Efficacy and safety of topical nadifloxacin and benzoyl peroxide versus clindamycin and benzoyl peroxide in acne vulgaris: A randomized controlled trial. *Indian J Pharmacol.* 2011;43; 628-631.
- [23] Christian GL, Krueger GG. Clindamycin vs placebo as adjunctive therapy in moderately severe acne. *Arch Dermatol.* 1975;111; 997-1000.
- [24] Christiansen J, Holm P, Reymann F. The retinoic acid derivative Ro 11-1430 in Acne vulgaris. A controlled multicenter trial against retinoic acid. *Dermatologica.* 1977;154; 219-227.
- [25] Chu A, Huber FJ, Plott RT. The comparative efficacy of benzoyl peroxide 5%/erythromycin 3% gel and erythromycin 4%/zinc 1.2% solution in the treatment of acne vulgaris. *Br J Dermatol.* 1997;136; 235-238.
- [26] Cullen SI, Cohan RH. Minocycline therapy in acne vulgaris. *Cutis.* 1976;17; 1208-1210, 1214.
- [27] Cunliffe WJ, Burke B, Dodman B, Gould DJ. A double-blind trial of a zinc sulphate/citrate complex and tetracycline in the treatment of acne vulgaris. *Br J Dermatol.* 1979;101; 321-325.
- [28] Cunliffe WJ, Caputo R, Dreno B, Forstrom L, Heenen M, Orfanos CE, et al. Efficacy and safety comparison of adapalene (CD271) gel and tretinoin gel in the topical treatment of acne vulgaris. A European multicentre trial. *Journal of Dermatological Treatment.* 1997;8; 173-178.
- [29] Cunliffe WJ, Danby FW, Dunlap F, Gold MH, Gratton D, Greenspan A. Randomised, controlled trial of the efficacy and safety of adapalene gel 0.1% and tretinoin cream 0.05% in patients with acne vulgaris. *Eur J Dermatol.* 2002;12; 350-354.
- [30] Cunliffe WJ, Fernandez C, Bojar R, Kanis R, West F. An observer-blind parallel-group, randomized, multicentre clinical and microbiological study of a topical clindamycin/zinc gel and a topical clindamycin lotion in patients with mild/moderate acne. *The Journal of dermatological treatment.* 2005;16; 213-218.
- [31] Cunliffe WJ, Grosshans E, Belaich S, Meynadier J, Alirezai M, Thomas L. A comparison of the efficacy and safety of lymecycline and minocycline in patients with moderately severe acne vulgaris. *Eur J Dermatol.* 1998;8; 161-166.
- [32] Cunliffe WJ, Holland KT. Clinical and laboratory studies on treatment with 20% azelaic acid cream for acne. *Acta Derm Venereol Suppl (Stockh).* 1989;143; 31-34.
- [33] Cunliffe WJ, Holland KT, Bojar R, Levy SF. A randomized, double-blind comparison of a clindamycin phosphate/benzoyl peroxide gel formulation and a matching clindamycin gel with respect to microbiologic activity and clinical efficacy in the topical treatment of acne vulgaris. *Clin Ther.* 2002;24; 1117-1133.
- [34] Cunliffe WJ, Meynadier J, Alirezai M, George SA, Coutts I, Roseeuw DI, et al. Is combined oral and topical therapy better than oral therapy alone in patients with moderate to moderately severe acne vulgaris? A comparison of the efficacy and

- safety of lymecycline plus adapalene gel 0.1%, versus lymecycline plus gel vehicle. *J Am Acad Dermatol.* 2003;49; S218-226.
- [35] Del Rosso J. Study results of benzoyl peroxide 5%/clindamycin J Drugs Dermatol. 2007;6; 616-622.
- [36] Dhawan SS. Comparison of 2 clindamycin 1%-benzoyl peroxide 5% topical gels used once daily in the management of acne vulgaris. *Cutis.* 2009;83; 265-272.
- [37] do Nascimento LV, Guedes AC, Magalhaes GM, de Faria FA, Guerra RM, de C. Almeida, F. Single-blind and comparative clinical study of the efficacy and safety of benzoyl peroxide 4% gel (BID) and adapalene 0.1% Gel (QD) in the treatment of acne vulgaris for 11 weeks. *The Journal of dermatological treatment.* 2003;14; 166-171.
- [38] Dobson RL, Belknap BS. Topical erythromycin solution in acne. Results of a multiclinic trial. *J Am Acad Dermatol.* 1980;3; 478-482.
- [39] Dominguez J, Hojyo MT, Celayo JL, Dominguez-Soto L, Teixeira F. Topical isotretinoin vs. topical retinoic acid in the treatment of acne vulgaris. *Int J Dermatol.* 1998;37; 54-55.
- [40] Draelos ZD, Shalita AR, Thiboutot D, Oresajo C, Yatskayer M, Raab S. A multicenter, double-blind study to evaluate the efficacy and safety of 2 treatments in participants with mild to moderate acne vulgaris. *Cutis.* 2012;89; 287-293.
- [41] Dreno B, Amblard P, Agache P, Sirot S, Litoux P. Low doses of zinc gluconate for inflammatory acne. *Acta Derm Venereol.* 1989;69; 541-543.
- [42] Dreno B, Kaufmann R, Talarico S, Torres Lozada V, Rodriguez-Castellanos MA, Gomez-Flores M, et al. Combination therapy with adapalene-benzoyl peroxide and oral lymecycline in the treatment of moderate to severe acne vulgaris: a multicentre, randomized, double-blind controlled study. *Br J Dermatol.* 2011;165; 383-390.
- [43] Dreno B, Moyse D, Alirezai M, Amblard P, Auffret N, Beylot C, et al. Multicenter randomized comparative double-blind controlled clinical trial of the safety and efficacy of zinc gluconate versus minocycline hydrochloride in the treatment of inflammatory acne vulgaris. *Dermatology.* 2001;203; 135-140.
- [44] Dubertret L, Alirezai M, Rostain G, Lahfa M, Forsea D, Niculae BD, et al. The use of lymecycline in the treatment of moderate to severe acne vulgaris: a comparison of the efficacy and safety of two dosing regimens. *Eur J Dermatol.* 2003;13; 44-48.
- [45] Ede M. A double-blind, comparative study of benzoyl peroxide, benzoyl peroxide-chlorhydroxyquinoline, benzoyl peroxide-chlorhydroxyquinoline-hydrocortisone, and placebo lotions in acne. *Curr Ther Res Clin Exp.* 1973;15; 624-629.
- [46] Eichenfield LF, Alio Saenz AB. Safety and efficacy of clindamycin phosphate 1.2%-benzoyl peroxide 3% fixed-dose combination gel for the treatment of acne vulgaris: a phase 3, multicenter, randomized, double-blind, active- and vehicle-controlled study. *J Drugs Dermatol.* 2011;10; 1382-1396.
- [47] Eichenfield LF, Draelos Z, Lucky AW, Hebert AA, Sugarman J, Stein Gold L, et al. Preadolescent moderate acne vulgaris: a randomized trial of the efficacy and safety of topical adapalene-benzoyl peroxides. *J Drugs Dermatol.* 2013;12; 611-618.
- [48] Eichenfield LF, Hebert AA, Schachner L, Paller AS, Rossi AB, Lucky AW. Tretinoin microsphere gel 0.04% pump for treating acne vulgaris in preadolescents: a randomized, controlled study. *Pediatr Dermatol.* 2012;29; 598-604.
- [49] Eichenfield LF, Jarratt M, Schlessinger J, Kempers S, Manna V, Hwa J, et al. Adapalene 0.1% lotion in the treatment of acne vulgaris: results from two placebo-controlled, multicenter, randomized double-blind, clinical studies. *J Drugs Dermatol.* 2010;9; 639-646.
- [50] Ellis CN, Gammon WR, Stone DZ, Heezen-Wehner JL. A comparison of Cleocin T Solution, Cleocin T Gel, and placebo in the treatment of acne vulgaris. *Cutis.* 1988;42; 245-247.
- [51] Ellis CN, Millikan LE, Smith EB, Chalker DM, Swinyer LJ, Katz IH, et al. Comparison of adapalene 0.1% solution and tretinoin 0.025% gel in the topical treatment of acne vulgaris. *Br J Dermatol.* 1998;139 Suppl 52; 41-47.

- [52] Fagundes DS, Fraser JM, Klauda HC. New therapy update--A unique combination formulation in the treatment of inflammatory acne. *Cutis*. 2003;72; 16-19.
- [53] Feucht CL, Allen BS, Chalker DK, Smith JG, Jr. Topical erythromycin with zinc in acne. A double-blind controlled study. *J Am Acad Dermatol*. 1980;3; 483-491.
- [54] Fleischer AB, Jr., Dinehart S, Stough D, Plott RT. Safety and efficacy of a new extended-release formulation of minocycline. *Cutis*. 2006;78; 21-31.
- [55] Fyrand O, Jakobsen HB. Water-based versus alcohol-based benzoyl peroxide preparations in the treatment of acne vulgaris. *Dermatologica*. 1986;172; 263-267.
- [56] Gammon WR, Meyer C, Lantis S, Shenefelt P, Reizner G, Cripps DJ. Comparative efficacy of oral erythromycin versus oral tetracycline in the treatment of acne vulgaris. A double-blind study. *J Am Acad Dermatol*. 1986;14; 183-186.
- [57] Glass D, Boorman GC, Stables GI, Cunliffe WJ, Goode K. A placebo-controlled clinical trial to compare a gel containing a combination of isotretinoin (0.05%) and erythromycin (2%) with gels containing isotretinoin (0.05%) or erythromycin (2%) alone in the topical treatment of acne vulgaris. *Dermatology*. 1999;199; 242-247.
- [58] Gollnick HP, Draehos Z, Glenn MJ, Rosoph LA, Kaszuba A, Cornelison R, et al. Adapalene-benzoyl peroxide, a unique fixed-dose combination topical gel for the treatment of acne vulgaris: a transatlantic, randomized, double-blind, controlled study in 1670 patients. *Br J Dermatol*. 2009;161; 1180-1189.
- [59] Gollnick HP, Graupe K, Zaumseil RP. Comparison of combined azelaic acid cream plus oral minocycline with oral isotretinoin in severe acne. *Eur J Dermatol*. 2001;11; 538-544.
- [60] Gollnick HP, Graupe K, Zaumseil RP. [Azelaic acid 15% gel in the treatment of acne vulgaris. Combined results of two double-blind clinical comparative studies]. *J Dtsch Dermatol Ges*. 2004;2; 841-847.
- [61] Goltz RW, Coryell GM, Schnieders JR, Neidert GL. A comparison of Cleocin T 1 percent solution and Cleocin T 1 percent lotion in the treatment of acne vulgaris. *Cutis*. 1985;36; 265-268.
- [62] Gonzalez P, Vila R, Cirigliano M. The tolerability profile of clindamycin 1%/benzoyl peroxide 5% gel vs. adapalene 0.1%/benzoyl peroxide 2.5% gel for facial acne: results of a randomized, single-blind, split-face study. *J*. 2012;11; 251-260.
- [63] Gratton D, Raymond GP, Guertin-Larochelle S, Maddin SW, Leneck CM, Warner J, et al. Topical clindamycin versus systemic tetracycline in the treatment of acne. Results of a multiclinic trial. *J Am Acad Dermatol*. 1982;7; 50-53.
- [64] Grosshans E, Marks R, Mascaro JM, Torras H, Meynadier J, Alirezai M, et al. Evaluation of clinical efficacy and safety of adapalene 0.1% gel versus tretinoin 0.025% gel in the treatment of acne vulgaris, with particular reference to the onset of action and impact on quality of life. *Br J Dermatol*. 1998;139 Suppl 52; 26-33.
- [65] Guerra-Tapia A. Effects of benzoyl peroxide 5% clindamycin combination gel versus adapalene 0.1% on quality of life in patients with mild to moderate acne vulgaris: a randomized single-blind study. *J Drugs Dermatol*. 2012;11; 714-722.
- [66] Gupta AK, Lynde CW, Kunynetz RA, Amin S, Choi K, Goldstein E. A randomized, double-blind, multicenter, parallel group study to compare relative efficacies of the topical gels 3% erythromycin/5% benzoyl peroxide and 0.025% tretinoin/erythromycin 4% in the treatment of moderate acne vulgaris of the face. *J Cutan Med Surg*. 2003;7; 31-37.
- [67] Habbema L, Koopmans B, Menke HE, Doornweerd S, De Boulle K. A 4% erythromycin and zinc combination (Zineryt) versus 2% erythromycin (Eryderm) in acne vulgaris: a randomized, double-blind comparative study. *Br J Dermatol*. 1989;121; 497-502.
- [68] Handojo I. Retinoic acid cream (Airol cream) and benzoyl-peroxide in the treatment of acne vulgaris. *Southeast Asian J Trop Med Public Health*. 1979;10; 548-551.
- [69] Harrison PV. A comparison of doxycycline and minocycline in the treatment of acne vulgaris. *Clin Exp Dermatol*. 1988;13; 242-244.
- [70] Hayashi N, Koyanagi E, Nogita T, Fujiyama M, Kawashima M. A randomized placebo-controlled investigator-blinded face split study of 20% azelaic acid cream to

- evaluate the efficacy and safety in Japanese patients with acne vulgaris. *J Dermatol.* 2012;39; 249-250.
- [71] Hughes BR, Norris JF, Cunliffe WJ. A double-blind evaluation of topical isotretinoin 0.05%, benzoyl peroxide gel 5% and placebo in patients with acne. *Clin Exp Dermatol.* 1992;17; 165-168.
- [72] Hunt MJ, Barnetson RS. A comparative study of gluconolactone versus benzoyl peroxide in the treatment of acne. *Australas J Dermatol.* 1992;33; 131-134.
- [73] Ioannides D, Rigopoulos D, Katsambas A. Topical adapalene gel 0.1% vs. isotretinoin gel 0.05% in the treatment of acne vulgaris: a randomized open-label clinical trial. *Br J Dermatol.* 2002;147; 523-527.
- [74] Iraji F, Sadeghinia A, Shahmoradi Z, Siadat AH, Jooya A. Efficacy of topical azelaic acid gel in the treatment of mild-moderate acne vulgaris. *Indian J Dermatol Venereol Leprol.* 2007;73; 94-96.
- [75] Jackson JM, Fu JJ, Almekinder JL. A randomized, investigator-blinded trial to assess the antimicrobial efficacy of a benzoyl peroxide 5%/ clindamycin phosphate 1% gel compared with a clindamycin phosphate 1.2%/tretinoin 0.025% gel in the topical treatment of acne vulgaris. *J Drugs Dermatol.* 2010;9; 131-136.
- [76] Jarratt MT, Brundage T. Efficacy and safety of clindamycin-tretinoin gel versus clindamycin or tretinoin alone in acne vulgaris: a randomized, double-blind, vehicle-controlled study. *J Drugs Dermatol.* 2012;11; 318-326.
- [77] Jones DH, King K, Miller AJ, Cunliffe WJ. A dose-response study of 13-cis-retinoic acid in acne vulgaris. *Br J Dermatol.* 1983;108; 333-343.
- [78] Jones EL, Crumley AF. Topical erythromycin vs blank vehicle in a multiclinic acne study. *Arch Dermatol.* 1981;117; 551-553.
- [79] Jung JY, Kwon HH, Yeom KB, Yoon MY, Suh DH. Clinical and histological evaluation of 1% nadifloxacin cream in the treatment of acne vulgaris in Korean patients. *Int J Dermatol.* 2011;50; 350-357.
- [80] Katsambas A, Graupe K, Stratigos J. Clinical studies of 20% azelaic acid cream in the treatment of acne vulgaris. Comparison with vehicle and topical tretinoin. *Acta Derm Venereol Suppl (Stockh).* 1989;143; 35-39.
- [81] Katsambas A, Towarky AA, Stratigos J. Topical clindamycin phosphate compared with oral tetracycline in the treatment of acne vulgaris. *Br J Dermatol.* 1987;116; 387-391.
- [82] Kaur J, Sehgal VK, Gupta AK, Singh SP. A comparative study to evaluate the efficacy and safety of combination topical preparations in acne vulgaris. *International journal of applied & basic medical research.* 2015;5; 106-110.
- [83] Kawashima M, Harada S, Loesche C, Miyachi Y. Adapalene gel 0.1% is effective and safe for Japanese patients with acne vulgaris: a randomized, multicenter, investigator-blinded, controlled study. *J Dermatol Sci.* 2008;49; 241-248.
- [84] Kawashima M, Hashimoto H, Alio Saenz AB, Ono M, Yamada M. Is benzoyl peroxide 3% topical gel effective and safe in the treatment of acne vulgaris in Japanese patients? A multicenter, randomized, double-blind, vehicle-controlled, parallel-group study. *J Dermatol.* 2014;41; 795-801.
- [85] Kawashima M, Hashimoto H, Saenz ABA, Ono M, Yamada M. Clindamycin phosphate 1.2%-benzoyl peroxide 3.0% fixed-dose combination gel has an effective and acceptable safety and tolerability profile for the treatment of acne vulgaris in Japanese patients: a phase III, multicentre, randomised, single-blinded, active-controlled, parallel-group study. *Br J Dermatol.* 2015;172; 494-503.
- [86] Khodaeinai E, Babaeinejad S, Amirnia M, Shokry J, Karimi LR, Fouladi DF, et al. Efficacy of 10% azelaic acid gel with hydro-alcoholic or alcohol-free bases in mild to moderate acne vulgaris; the first clinical trial. *Journal of Medical Sciences (Faisalabad).* 2014;14; 87-91.
- [87] Kircik L. Community-based trial results of combination clindamycin 1%--benzoyl peroxide 5% topical gel plus tretinoin microsphere gel 0.04% or 0.1% or adapalene gel 0.1% in the treatment of moderate to severe acne. *Cutis.* 2007;80; 10-14.

- [88] Kircik L, Green L, Thiboutot D, Tanghetti E, Wilson D, Dhawan S, et al. Comparing a novel solubilized benzoyl peroxide gel with benzoyl peroxide/clindamycin: final data from a multicenter, investigator-blind, randomized study. *J Drugs Dermatol.* 2009;8; 812-818.
- [89] Kircik LH. Comparative efficacy and safety results of two topical combination acne regimens. *J Drugs Dermatol.* 2009;8; 624-630.
- [90] Ko HC, Song M, Seo SH, Oh CK, Kwon KS, Kim MB. Prospective, open-label, comparative study of clindamycin 1%/benzoyl peroxide 5% gel with adapalene 0.1% gel in Asian acne patients: efficacy and tolerability. *J Eur Acad Dermatol Venereol.* 2009;23; 245-250.
- [91] Kobayashi M, Nakagawa T, Fukamachi K, Nakamura M, Tokura Y. Efficacy of combined topical treatment of acne vulgaris with adapalene and nadifloxacin: a randomized study. *J Dermatol.* 2011;38; 1163-1166.
- [92] Korkut C, Piskin S. Benzoyl peroxide, adapalene, and their combination in the treatment of acne vulgaris. *J Dermatol.* 2005;32; 169-173.
- [93] Kuhlman DS, Callen JP. A comparison of clindamycin phosphate 1 percent topical lotion and placebo in the treatment of acne vulgaris. *Cutis.* 1986;38; 203-206.
- [94] Langner A, Boorman GC, Stapor V. Isotretinoin cream 0.05% and 0.1% in the treatment of acne vulgaris. *J Dermatol Treat.* 1994;5; 177-180.
- [95] Langner A, Chu A, Goulden V, Ambroziak M. A randomized, single-blind comparison of topical clindamycin + benzoyl peroxide and adapalene in the treatment of mild to moderate facial acne vulgaris. *Br J Dermatol.* 2008;158; 122-129.
- [96] Langner A, Sheehan-Dare R, Layton A. A randomized, single-blind comparison of topical clindamycin + benzoyl peroxide (Duac) and erythromycin + zinc acetate (Zineryt) in the treatment of mild to moderate facial acne vulgaris. *J Eur Acad Dermatol Venereol.* 2007;21; 311-319.
- [97] Lee JW, Yoo KH, Park KY, Han TY, Li K, Seo SJ, et al. Effectiveness of conventional, low-dose and intermittent oral isotretinoin in the treatment of acne: a randomized, controlled comparative study. *Br J Dermatol.* 2011;164; 1369-1375.
- [98] Lesher JL, Jr., Chalker DK, Smith JG, Jr., Guenther LC, Ellis CN, Voorhees JJ, et al. An evaluation of a 2% erythromycin ointment in the topical therapy of acne vulgaris. *J Am Acad Dermatol.* 1985;12; 526-531.
- [99] Lester RS, Schachter GD, Light MJ. Isotretinoin and tetracycline in the management of severe nodulocystic acne. *Int J Dermatol.* 1985;24; 252-257.
- [100] Leyden JJ, Berger RS, Dunlap FE, Ellis CN, Connolly MA, Levy SF. Comparison of the efficacy and safety of a combination topical gel formulation of benzoyl peroxide and clindamycin with benzoyl peroxide, clindamycin and vehicle gel in the treatments of acne vulgaris. *American journal of clinical dermatology.* 2001;2; 33-39.
- [101] Leyden JJ, Bruce S, Lee CS, Ling M, Sheth PB, Stewart DM, et al. A randomized, phase 2, dose-ranging study in the treatment of moderate to severe inflammatory facial acne vulgaris with doxycycline calcium. *J Drugs Dermatol.* 2013;12; 658-663.
- [102] Leyden JJ, Hickman JG, Jarratt MT, Stewart DM, Levy SF. The efficacy and safety of a combination benzoyl peroxide/clindamycin topical gel compared with benzoyl peroxide alone and a benzoyl peroxide/erythromycin combination product. *J Cutan Med Surg.* 2001;5; 37-42.
- [103] Leyden JJ, Krochmal L, Yaroshinsky A. Two randomized, double-blind, controlled trials of 2219 subjects to compare the combination clindamycin/tretinoin hydrogel with each agent alone and vehicle for the treatment of acne vulgaris. *J Am Acad Dermatol.* 2006;54; 73-81.
- [104] Leyden JJ, Shalita AR, Saatjian GD, Sefton J. Erythromycin 2% gel in comparison with clindamycin phosphate 1% solution in acne vulgaris. *J Am Acad Dermatol.* 1987;16; 822-827.
- [105] Lookingbill DP, Chalker DK, Lindholm JS, Katz HI, Kempers SE, Huerter CJ, et al. Treatment of acne with a combination clindamycin/benzoyl peroxide gel compared with clindamycin gel, benzoyl peroxide gel and vehicle gel: combined results of two double-blind investigations. *J Am Acad Dermatol.* 1997;37; 590-595.

- [106] Lucchini LC, Kollias N, Gillies R, Phillips SB, Muccini JA, Stiller MJ, et al. Fluorescence photography in the evaluation of acne. *J Am Acad Dermatol.* 1996;35; 58-63.
- [107] Lucky A, Jorizzo JL, Rodriguez D, Jones TM, Stewart DM, Tschen EH, et al. Efficacy and tolerance of adapalene cream 0.1% compared with its cream vehicle for the treatment of acne vulgaris. *Cutis.* 2001;68; 34-40.
- [108] Lucky AW, Cullen SI, Funicella T, Jarratt MT, Jones T, Reddick ME. Double-blind, vehicle-controlled, multicenter comparison of two 0.025% tretinoin creams in patients with acne vulgaris. *J Am Acad Dermatol.* 1998;38; S24-30.
- [109] Lucky AW, Cullen SI, Jarratt MT, Quigley JW. Comparative efficacy and safety of two 0.025% tretinoin gels: results from a multicenter double-blind, parallel study. *J Am Acad Dermatol.* 1998;38; S17-23.
- [110] Lyons RE. Comparative effectiveness of benzoyl peroxide and tretinoin in acne vulgaris. *Int J Dermatol.* 1978;17; 246-251.
- [111] Marazzi P, Boorman GC, Donald AE, Davies HD. Clinical evaluation of Double Strength Isotrexin versus Benzamycin in the topical treatment of mild to moderate acne vulgaris. *The Journal of dermatological treatment.* 2002;13; 111-117.
- [112] McKenzie MW, Beck DC, Popovich NG. Topical clindamycin formulations for the treatment of acne vulgaris. An evaluation. *Arch Dermatol.* 1981;117; 630-634.
- [113] Mills O, Jr., Thornsberry C, Cardin CW, Smiles KA, Leyden JJ. Bacterial resistance and therapeutic outcome following three months of topical acne therapy with 2% erythromycin gel versus its vehicle. *Acta Derm Venereol.* 2002;82; 260-265.
- [114] Mills OH, Berger RS, Kligman AM, McElroy JA, Di Matteo J. A comparative study of Erycette(TM) vs Cleocin-T(TM). *Adv Ther.* 1992;9; 14-20.
- [115] Moore ALMBAMVRMJ. Efficacy and safety of subantimicrobial dose, modified-release doxycycline 40 mg versus doxycycline 100 mg versus placebo for the treatment of inflammatory lesions in Moderate and Severe Acne: A Randomized, Double-Blinded, Controlled Study. *Journal of Drugs in Dermatology.* 2015;14; 581-586.
- [116] Nandimath MK, Reddy NB. Comparision of clinical efficacy of topical clindamycin with adapalene and adapalene alone in treatment of mild to moderate facial acne vulgaris. *International Journal of Pharma and Bio Sciences.* 2013;4; P1079-P1088.
- [117] Nighland M, Grossman R. Tretinoin microsphere gel in facial acne vulgaris: a meta-analysis. *J Drugs Dermatol.* 2008;7; s2-8.
- [118] NilFroushzadeh MA, Siadat AH, Baradaran EH, Moradi S. Clindamycin lotion alone versus combination lotion of clindamycin phosphate plus tretinoin versus combination lotion of clindamycin phosphate plus salicylic acid in the topical treatment of mild to moderate acne vulgaris: a randomized control trial. *Indian J Dermatol Venereol Leprol.* 2009;75; 279-282.
- [119] Norris JF, Hughes BR, Basey AJ, Cunliffe WJ. A comparison of the effectiveness of topical tetracycline, benzoyl-peroxide gel and oral oxytetracycline in the treatment of acne. *Clin Exp Dermatol.* 1991;16; 31-33.
- [120] Nyirady J, Grossman RM, Nighland M, Berger RS, Jorizzo JL, Kim YH, et al. A comparative trial of two retinoids commonly used in the treatment of acne vulgaris. *The Journal of dermatological treatment.* 2001;12; 149-157.
- [121] Ocampo-Candiani J, Velazquez-Arenas LL, de la Fuente-Garcia A, Trevino-Gomezharper C, Berber A. Safety and efficacy comparison of minocycline microgranules vs lymecycline in the treatment of mild to moderate acne: randomized, evaluator-blinded, parallel, and prospective clinical trial for 8 weeks. *J Drugs Dermatol.* 2014;13; 671-676.
- [122] Olafsson JH, Gudgerisson J, Eggertsottir GE, Kristjansson F. Doxycycline versus minocycline in the treatment of acne vulgaris: A double-blind study. *Journal of Dermatological Treatment.* 1989;1; 15-17.
- [123] Oprica C, Emtestam L, Hagstromer L, Nord CE. Clinical and microbiological comparisons of isotretinoin vs. tetracycline in acne vulgaris. *Acta Derm Venereol.* 2007;87; 246-254.

- [124] Orris L, Shalita AR, Sibulkin D, London SJ, Gans EH. Oral zinc therapy of acne. Absorption and clinical effect. *Arch Dermatol.* 1978;114; 1018-1020.
- [125] Ozgen ZY, Gurbuz O. A randomized, double-blind comparison of nadifloxacin 1% cream alone and with benzoyl peroxide 5% lotion in the treatment of mild to moderate facial acne vulgaris
Hafif-orta sidetli akne vulgaris yuz tutulumu tedavisinde nadifloksasinin tek basi{dotless}na ve benzoil peroksit ile birlikte kullan{dotless}mi{dotless}ni{dotless}n karsi{dotless}lasti{dotless}ri{dotless}lmasi{dotless}. Marmara Medical Journal. 2013;26.
- [126] Ozolins M, Eady EA, Avery A, Cunliffe WJ, O'Neill C, Simpson NB, et al. Randomised controlled multiple treatment comparison to provide a cost-effectiveness rationale for the selection of antimicrobial therapy in acne. *Health Technol Assess.* 2005;9; iii-212.
- [127] Ozolins M, Eady EA, Avery AJ, Cunliffe WJ, Po AL, O'Neill C, et al. Comparison of five antimicrobial regimens for treatment of mild to moderate inflammatory facial acne vulgaris in the community: randomised controlled trial. *Lancet.* 2004;364; 2188-2195.
- [128] Packman AM, Brown RH, Dunlap FE, Kraus SJ, Webster GF. Treatment of acne vulgaris: combination of 3% erythromycin and 5% benzoyl peroxide in a gel compared to clindamycin phosphate lotion. *Int J Dermatol.* 1996;35; 209-211.
- [129] Padilla RS, McCabe JM, Becker LE. Topical tetracycline hydrochloride vs. topical clindamycin phosphate in the treatment of acne: a comparative study. *Int J Dermatol.* 1981;20; 445-448.
- [130] Panzer JD, Poche W, Meek TJ, Derbes VJ, Atkinson W. Acne treatment: a comparative efficacy trial of clindamycin and tetracycline. *Cutis.* 1977;19; 109-111.
- [131] Papageorgiou PP, Chu AC. Chloroxylenol and zinc oxide containing cream (Nels cream) vs. 5% benzoyl peroxide cream in the treatment of acne vulgaris. A double-blind, randomized, controlled trial. *Clin Exp Dermatol.* 2000;25; 16-20.
- [132] Pariser D, Bucko A, Fried R, Jarratt MT, Kempers S, Kircik L, et al. Tretinoin gel microsphere pump 0.04% plus 5% benzoyl peroxide wash for treatment of acne vulgaris: morning/morning regimen is as effective and safe as morning/evening regimen. *J Drugs Dermatol.* 2010;9; 805-813.
- [133] Pariser DM, Rich P, Cook-Bolden FE, Korotzer A. An aqueous gel fixed combination of clindamycin phosphate 1.2% and benzoyl peroxide 3.75% for the once-daily treatment of moderate to severe acne vulgaris. *J Drugs Dermatol.* 2014;13; 1083-1089.
- [134] Pariser DM, Thiboutot DM, Clark SD, Jones TM, Liu Y, Graeber M. The efficacy and safety of adapalene gel 0.3% in the treatment of acne vulgaris: A randomized, multicenter, investigator-blinded, controlled comparison study versus adapalene gel 0.1% and vehicle. *Cutis.* 2005;76; 145-151.
- [135] Parker F. A comparison of clindamycin 1% solution versus clindamycin 1% gel in the treatment of acne vulgaris. *Int J Dermatol.* 1987;26; 121-122.
- [136] Pazoki-Toroudi H, Nassiri-Kashani M, Tabatabaie H, Ajami M, Habibey R, Shizarpour M, et al. Combination of azelaic acid 5% and erythromycin 2% in the treatment of acne vulgaris. *J Dermatolog Treat.* 2010;21; 212-216.
- [137] Pazoki-Toroudi H, Nilforoushzadeh MA, Ajami M, Jaffary F, Aboutaleb N, Nassiri-Kashani M, et al. Combination of azelaic acid 5% and clindamycin 2% for the treatment of acne vulgaris. *Cutan.* 2011;30; 286-291.
- [138] Peacock CE, Price C, Ryan BE, Mitchell AD. Topical clindamycin (Dalaclin T(TM)) compared to oral minocycline (Minocin 50(TM)) in treatment of acne vulgaris. A randomized observer-blind controlled trial in three university student health centres. *Clin Trials J.* 1990;27; 219-228.
- [139] Peck GL, Olsen TG, Butkus D, Pandya M, Arnaud-Battandier J, Gross EG, et al. Isotretinoin versus placebo in the treatment of cystic acne. A randomized double-blind study. *J Am Acad Dermatol.* 1982;6; 735-745.
- [140] Petersen MJ, Krusinski PA, Krueger GG. Evaluation of 1% clindamycin phosphate lotion in the treatment of acne. *Curr Ther Res.* 1986;40; 232-238.

- [141] Pierard-Franchimont C, Goffin V, Arrese JE, Martalo O, Braham C, Slachmuylders P, et al. Lymecycline and minocycline in inflammatory acne: a randomized, double-blind intent-to-treat study on clinical and in vivo antibacterial efficacy. *Skin Pharmacol Appl Skin Physiol.* 2002;15; 112-119.
- [142] Pierard-Franchimont C, Henry F, Fraiture AL, Fumal I, Pierard GE. Split-face clinical and bio-instrumental comparison of 0.1% adapalene and 0.05% tretinoin in facial acne. *Dermatology.* 1999;198; 218-222.
- [143] Pigatto PD, Finzi AF, Altomare GF, Polenghi MM, Vergani C, Vigotti G. Isotretinoin versus minocycline in cystic acne: a study of lipid metabolism. *Dermatologica.* 1986;172; 154-159.
- [144] Plewig G, Holland KT, Nenoff P. Clinical and bacteriological evaluation of nadifloxacin 1% cream in patients with acne vulgaris: a double-blind, phase III comparison study versus erythromycin 2% cream. *Eur J Dermatol.* 2006;16; 48-55.
- [145] Pochi PE, Bagatell FK, Ellis CN, Stoughton RB, Whitmore CG, Saatjian GD, et al. Erythromycin 2 percent gel in the treatment of acne vulgaris. *Cutis.* 1988;41; 132-136.
- [146] Prasad S, Mukhopadhyay A, Kubavat A, Kelkar A, Modi A, Swarnkar B, et al. Efficacy and safety of a nano-emulsion gel formulation of adapalene 0.1% and clindamycin 1% combination in acne vulgaris: a randomized, open label, active-controlled, multicentric, phase IV clinical trial. *Indian J Dermatol Venereol Leprol.* 2012;78; 459-467.
- [147] Rademaker M, Wishart JM, Birchall NM. Isotretinoin 5 mg daily for low-grade adult acne vulgaris--a placebo-controlled, randomized double-blind study. *J Eur Acad Dermatol Venereol.* 2014;28; 747-754.
- [148] Rapaport M, Puhvel SM, Reisner RM. Evaluation of topical erythromycin and oral tetracycline in acne vulgaris. *Cutis.* 1982;30; 122-126, 130, 132-125.
- [149] Richter JR, Forstrom LR, Kiistala UO, Jung EG. Efficacy of the fixed 1.2% clindamycin phosphate, 0.025% tretinoin gel formulation (Velac) and a proprietary 0.025% tretinoin gel formulation (Aberela) in the topical control of facial acne. *J Eur Acad Dermatol Venereol.* 1998;11; 227-233.
- [150] Rietschel RL, Duncan SH. Clindamycin phosphate used in combination with tretinoin in the treatment of acne. *Int J Dermatol.* 1983;22; 41-43.
- [151] Rivkin L, Rapaport M. Clinical evaluation of a new erythromycin solution for acne vulgaris. *Cutis.* 1980;25; 552-555.
- [152] Rizer RL, Sklar JL, Whiting D, Bucko A, Shavin J, Jarratt M. Clindamycin phosphate 1% gel in acne vulgaris. *Adv Ther.* 2001;18; 244-252.
- [153] Robledo AA, Lopez BE, del Pino GJ, Sambricio GF, Rodriguez PA, Sotillo G, et al. Multicentric comparative study of the efficacy and tolerance of clindamycin phosphate 1% topical solution and tetracycline topical solution for the treatment of acne vulgaris. *Curr Ther Res Clin Exp.* 1988;43; 21-26.
- [154] Ruping KW, Tronnier H. Acne therapy: Results of a multicentre study with minocycline. Royal Society of Medicine Services International Congress and Symposium Series. 1985;95; 109-119.
- [155] Schachner L, Eaglstein W, Kittles C, Mertz P. Topical erythromycin and zinc therapy for acne. *J Am Acad Dermatol.* 1990;22; 253-260.
- [156] Schachner L, Pestana A, Kittles C. A clinical trial comparing the safety and efficacy of a topical erythromycin-zinc formulation with a topical clindamycin formulation. *J Am Acad Dermatol.* 1990;22; 489-495.
- [157] Schafer-Korting M, Korting HC, Ponce-Poschl E. Liposomal tretinoin for uncomplicated acne vulgaris. *Clin Investig.* 1994;72; 1086-1091.
- [158] Schmidt N, Gans EH. Clindamycin 1.2% Tretinoin 0.025% Gel versus Clindamycin Gel Treatment in Acne Patients: A Focus on Fitzpatrick Skin Types. *J Clin Aesthet Dermatol.* 2011;4; 31-40.
- [159] Shahlita AR, Smith EB, Bauer E. Topical erythromycin v clindamycin therapy for acne. A multicenter, double-blind comparison. *Arch Dermatol.* 1984;120; 351-355.

- [160] Shalita A, Weiss JS, Chalker DK, Ellis CN, Greenspan A, Katz HI, et al. A comparison of the efficacy and safety of adapalene gel 0.1% and tretinoin gel 0.025% in the treatment of acne vulgaris: a multicenter trial. *J Am Acad Dermatol.* 1996;34; 482-485.
- [161] Shalita AR, Myers JA, Krochmal L, Yaroshinsky A. The safety and efficacy of clindamycin phosphate foam 1% versus clindamycin phosphate topical gel 1% for the treatment of acne vulgaris. *J Drugs Dermatol.* 2005;4; 48-56.
- [162] Shalita AR, Rafal ES, Anderson DN, Yavel R, Landow S, Lee WL. Compared efficacy and safety of tretinoin 0.1% microsphere gel alone and in combination with benzoyl peroxide 6% cleanser for the treatment of acne vulgaris. *Cutis.* 2003;72; 167-172.
- [163] Sheehan-Dare RA, Papworth-Smith J, Cunliffe WJ. A double-blind comparison of topical clindamycin and oral minocycline in the treatment of acne vulgaris. *Acta Derm Venereol.* 1990;70; 534-537.
- [164] Shwetha H, Geetha A, Revathi TN. A comparative study of efficacy and safety of combination of topical 1% clindamycin and 0.1% adapalene with 1% clindamycin and 2.5% benzoyl peroxide in mild to moderate acne at a tertiary care hospital. *Journal of Chemical and Pharmaceutical Research.* 2014;6; 736-741.
- [165] Skidmore R, Kovach R, Walker C, Thomas J, Bradshaw M, Leyden J, et al. Effects of subantimicrobial-dose doxycycline in the treatment of moderate acne. *Arch Dermatol.* 2003;139; 459-464.
- [166] Sklar JL, Jacobson C, Rizer R, Gans EH. Evaluation of Triaz 10% Gel and Benzamycin in acne vulgaris. *Journal of Dermatological Treatment.* 1996;7; 147-152.
- [167] Stainforth J, Macdonald-Hull S, Papworth-Smith JW, Eady EA, Cunliffe WJ, Norris JFB, et al. A single-blind comparison of topical erythromycin/zinc lotion and oral minocycline in the treatment of acne vulgaris. *Journal of Dermatological Treatment.* 1993;4; 119-122.
- [168] Stein Gold L, Cruz A, Eichenfield L, Tan J, Jorizzo J, Kerrouche N, et al. Effective and safe combination therapy for severe acne vulgaris: a randomized, vehicle-controlled, double-blind study of adapalene 0.1%-benzoyl peroxide 2.5% fixed-dose combination gel with doxycycline hydiate 100 mg. *Cutis.* 2010;85; 94-104.
- [169] Stein Gold L, Tan J, Cruz-Santana A, Papp K, Poulin Y, Schlessinger J, et al. A North American study of adapalene-benzoyl peroxide combination gel in the treatment of acne. *Cutis.* 2009;84; 110-116.
- [170] Stewart DM, Torok HM, Weiss JS, Plott RT. Dose-ranging efficacy of new once-daily extended-release minocycline for acne vulgaris. *Cutis.* 2006;78; 11-20.
- [171] Stinco G, Bragadin G, Trotter D, Pillon B, Patrone P. Relationship between sebostatic activity, tolerability and efficacy of three topical drugs to treat mild to moderate acne. *J Eur Acad Dermatol Venereol.* 2007;21; 320-325.
- [172] Strauss JS, Leyden JJ, Lucky AW, Lookingbill DP, Drake LA, Hanifin JM, et al. A randomized trial of the efficacy of a new micronized formulation versus a standard formulation of isotretinoin in patients with severe recalcitrant nodular acne. *J Am Acad Dermatol.* 2001;45; 187-195.
- [173] Strauss JS, Rapini RP, Shalita AR, Konecky E, Pochi PE, Comite H, et al. Isotretinoin therapy for acne: results of a multicenter dose-response study. *J Am Acad Dermatol.* 1984;10; 490-496.
- [174] Strauss JS, Stranieri AM. Acne treatment with topical erythromycin and zinc: effect of Propionibacterium acnes and free fatty acid composition. *J Am Acad Dermatol.* 1984;11; 86-89.
- [175] Swinyer LJ, Baker MD, Swinyer TA, Mills OH, Jr. A comparative study of benzoyl peroxide and clindamycin phosphate for treating acne vulgaris. *Br J Dermatol.* 1988;119; 615-622.
- [176] Takigawa M, Tokura Y, Shimada S, Furukawa F, Noguchi N, Ito T, et al. Clinical and bacteriological evaluation of adapalene 0.1% gel plus nadifloxacin 1% cream versus adapalene 0.1% gel in patients with acne vulgaris. *J Dermatol.* 2013;40; 620-625.
- [177] Tan J, Humphrey S, Vender R, Barankin B, Gooderham M, Kerrouche N, et al. A treatment for severe nodular acne: a randomized investigator-blinded, controlled,

- noninferiority trial comparing fixed-dose adapalene/benzoyl peroxide plus doxycycline vs. oral isotretinoin. *Br J Dermatol.* 2014;171; 1508-1516.
- [178] Tanghetti E, Kircik L, Wilson D, Dhawan S. Solubilized benzoyl peroxide versus benzoyl peroxide/clindamycin in the treatment of moderate acne. *J Drugs Dermatol.* 2008;7; 534-538.
- [179] Thiboutot D, Gold MH, Jarratt MT, Kang S, Kaplan DL, Millikan L, et al. Randomized controlled trial of the tolerability, safety, and efficacy of adapalene gel 0.1% and tretinoin microsphere gel 0.1% for the treatment of acne vulgaris. *Cutis.* 2001;68; 10-19.
- [180] Thiboutot D, Jarratt M, Rich P, Rist T, Rodriguez D, Levy S. A randomized, parallel, vehicle-controlled comparison of two erythromycin/benzoyl peroxide preparations for acne vulgaris. *Clin Ther.* 2002;24; 773-785.
- [181] Thiboutot D, Pariser DM, Egan N, Flores J, Herndon JH, Jr., Kanof NB, et al. Adapalene gel 0.3% for the treatment of acne vulgaris: a multicenter, randomized, double-blind, controlled, phase III trial. *J Am Acad Dermatol.* 2006;54; 242-250.
- [182] Thiboutot D, Zaenglein A, Weiss J, Webster G, Calvarese B, Chen D. An aqueous gel fixed combination of clindamycin phosphate 1.2% and benzoyl peroxide 2.5% for the once-daily treatment of moderate to severe acne vulgaris: assessment of efficacy and safety in 2813 patients. *J Am Acad Dermatol.* 2008;59; 792-800.
- [183] Thiboutot DM, Shalita AR, Yamauchi PS, Dawson C, Arsonnaud S, Kang S. Combination therapy with adapalene gel 0.1% and doxycycline for severe acne vulgaris: a multicenter, investigator-blind, randomized, controlled study. *Skinmed.* 2005;4; 138-146.
- [184] Thiboutot DM, Weiss J, Bucko A, Eichenfield L, Jones T, Clark S, et al. Adapalene-benzoyl peroxide, a fixed-dose combination for the treatment of acne vulgaris: results of a multicenter, randomized double-blind, controlled study. *J Am Acad Dermatol.* 2007;57; 791-799.
- [185] Thielitz A, Lux A, Wiede A, Kropf S, Papakonstantinou E, Gollnick H. A randomized investigator-blind parallel-group study to assess efficacy and safety of azelaic acid 15% gel vs. adapalene 0.1% gel in the treatment and maintenance treatment of female adult acne. *J Eur Acad Dermatol Venereol.* 2015;29; 789-796.
- [186] Thomas DR, Raimer S, Smith EB. Comparison of topical erythromycin 1.5 percent solution versus topical clindamycin phosphate 1.0 percent solution in the treatment of acne vulgaris. *Cutis.* 1982;29; 624-625, 628-632.
- [187] Tirado-Sanchez A, Espindola YS, Ponce-Olivera RM, Bonifaz A. Efficacy and safety of adapalene gel 0.1% and 0.3% and tretinoin gel 0.05% for acne vulgaris: results of a single-center, randomized, double-blinded, placebo-controlled clinical trial on Mexican patients (skin type III-IV). *J.* 2013;12; 103-107.
- [188] Toossi P, Farshchian M, Malekzad F, Mohtasham N, Kimyai-Asadi A. Subantimicrobial-dose doxycycline in the treatment of moderate facial acne. *J Drugs Dermatol.* 2008;7; 1149-1152.
- [189] Tschen EH, Katz HI, Jones TM, Monroe EW, Kraus SJ, Connolly MA, et al. A combination benzoyl peroxide and clindamycin topical gel compared with benzoyl peroxide, clindamycin phosphate, and vehicle in the treatment of acne vulgaris. *Cutis.* 2001;67; 165-169.
- [190] Tu P, Li GQ, Zhu XJ, Zheng J, Wong WZ. A comparison of adapalene gel 0.1% vs. tretinoin gel 0.025% in the treatment of acne vulgaris in China. *J Eur Acad Dermatol Venereol.* 2001;15 Suppl 3; 31-36.
- [191] Tucker SB, Tausend R, Cochran R, Flannigan SA. Comparison of topical clindamycin phosphate, benzoyl peroxide, and a combination of the two for the treatment of acne vulgaris. *Br J Dermatol.* 1984;110; 487-492.
- [192] Tunca M, Akar A, Ozmen I, Erbil H. Topical nadifloxacin 1% cream vs. topical erythromycin 4% gel in the treatment of mild to moderate acne. *Int J Dermatol.* 2010;49; 1440-1444.

- [193] Ustuner PGATDM. Clinical and bacteriological evaluation of nadifloxacin 1% cream versus erythromycin 4% gel in the treatment of mild-to-moderate facial acne vulgaris: A randomized study. *Turkiye Klinikleri Journal of Medical Sciences*. 2015;35; 97-105.
- [194] van der Meeren HL, van der Schroeff JG, Stijnen T, van Duren JA, van der Dries HA, van Voorst Vader PC. Dose-response relationship in isotretinoin therapy for conglobate acne. *Dermatologica*. 1983;167; 299-303.
- [195] Verschoore M, Langner A, Wolska H, Jablonska S, Czernielewski J, Schaefer H. Efficacy and safety of CD 271 alcoholic gels in the topical treatment of acne vulgaris. *Br J Dermatol*. 1991;124; 368-371.
- [196] Webster G, Cargill DL, Quiring J, Vogelson CT, Slade HB. A combined analysis of 2 randomized clinical studies of tretinoin gel 0.05% for the treatment of acne. *Cutis*. 2009;83; 146-154.
- [197] Webster G, Rich P, Gold MH, Mraz S, Calvarese B, Chen D. Efficacy and tolerability of a fixed combination of clindamycin phosphate (1.2%) and low concentration benzoyl peroxide (2.5%) aqueous gel in moderate or severe acne subpopulations. *J Drugs Dermatol*. 2009;8; 736-743.
- [198] Webster GF. Safety and efficacy of Tretin-X compared with Retin-A in patients with mild-to-severe acne vulgaris. *Skinmed*. 2006;5; 114-118.
- [199] Webster GF, Leyden JJ, Gross JA. Results of a Phase III, double-blind, randomized, parallel-group, non-inferiority study evaluating the safety and efficacy of isotretinoin-lidose in patients with severe recalcitrant nodular acne. *J Drugs Dermatol*. 2014;13; 665-670.
- [200] Weimar VM, Puhl SC, Smith WH, tenBroeke JE. Zinc sulfate in acne vulgaris. *Arch Dermatol*. 1978;114; 1776-1778.
- [201] Weismann K, Wadskov S, Sondergaard J. Oral zinc sulphate therapy for acne vulgaris. *Acta Derm Venereol*. 1977;57; 357-360.
- [202] Weiss JGLSTEBSYMDCLM. Efficacy and safety of adapalene 0.3%/benzoyl peroxide 2.5% topical gel in moderate and severe acne vulgaris. *J Am Acad Dermatol*. 2015;72; AB7.
- [203] Wolf JE, Jr., Kaplan D, Kraus SJ, Loven KH, Rist T, Swinyer LJ, et al. Efficacy and tolerability of combined topical treatment of acne vulgaris with adapalene and clindamycin: a multicenter, randomized, investigator-blinded study. *J Am Acad Dermatol*. 2003;49; S211-217.
- [204] Wong RC, Kang S, Heezen JL, Voorhees JJ, Ellis CN. Oral ibuprofen and tetracycline for the treatment of acne vulgaris. *J Am Acad Dermatol*. 1984;11; 1076-1081.
- [205] Yentzer BA, Ade RA, Fountain JM, Clark AR, Taylor SL, Fleischer AB, Jr., et al. Simplifying regimens promotes greater adherence and outcomes with topical acne medications: a randomized controlled trial. *Cutis*. 2010;86; 103-108.
- [206] Zhang JZ, Li LF, Tu YT, Zheng J. A successful maintenance approach in inflammatory acne with adapalene gel 0.1% after an initial treatment in combination with clindamycin topical solution 1% or after monotherapy with clindamycin topical solution 1%. *The Journal of dermatological treatment*. 2004;15; 372-378.
- [207] Zhu XJ, Tu P, Zhen J, Duan YQ. Adapalene gel 0.1%: effective and well tolerated in the topical treatment of acne vulgaris in Chinese patients. *Cutis*. 2001;68; 55-59.
- [208] Ziana [prescribing information]. Medicis , The Dermatology Company; Scottsdale, AZ.
- [209] Zouboulis CC, Derumeaux L, Decroix J, Maciejewska-Udziela B, Cambazard F, Stuhlert A. A multicentre, single-blind, randomized comparison of a fixed clindamycin phosphate/tretinoin gel formulation (Velac) applied once daily and a clindamycin lotion formulation (Dalacin T) applied twice daily in the topical treatment of acne vulgaris. *Br J Dermatol*. 2000;143; 498-505.
- [210] Zouboulis CC, Fischer TC, Wohlrab J, Barnard J, Alio AB. Study of the efficacy, tolerability, and safety of 2 fixed-dose combination gels in the management of acne vulgaris. *Cutis*. 2009;84; 223-229.